

IS THE DEMAND-CONTROL MODEL STILL A USEFUL TOOL TO ASSESS WORK-RELATED PSYCHOSOCIAL RISK FOR ISCHEMIC HEART DISEASE? RESULTS FROM 14 YEAR FOLLOW UP IN THE COPENHAGEN CITY HEART STUDY

BO NETTERSTRØM¹, TAGE S. KRISTENSEN², GORM JENSEN¹, and PETER SCHNOR¹

¹ Copenhagen City Heart Study, Bispebjerg Hospital, Copenhagen, Denmark

Department of Occupational and Environmental Medicine, Bispebjerg Hospital

² Task-Consult, Gilleleje, Denmark

Abstract

Objectives: To test the usefulness of the Demand-Control Model as predictor for ischemic heart disease (IHD).

Materials and Methods: One thousand one hundred forty six actively employed men and women from the general population of Copenhagen participated at baseline in 1993–1994. They filled in questionnaires on the Demand-Control Model, job title, work place, civil status, family income, leisure time activity, smoking, medication, social support, social relations, conflicts, job responsibility, satisfaction, and insecurity and went through a medical examination, including measurements of coronary risk factors. All deaths and hospital admissions due to IHD, including first myocardial infarction (MI) in the cohort were traced in the Danish registries of deaths and hospital admissions to June 2007. **Results:** 104 cases of first time hospitalisation or death due to IHD including 49 cases of MI occurred during 14 years follow up. Odds ratio (OR) compared to the relaxed group was 1.1 (0.1–3.1) among women and 1.6 (0.4–4.9) among men after confounder adjustment. Neither demands nor control were significantly associated with IHD. Among men 50 years of age or more, the risk for IHD was, however, elevated in the job strain group and the active group (OR = 3.5 and 3.2 respectively). Job insecurity was, however, strongly associated with IHD in men (OR = 2.7 (1.1–5.6)) after all adjustments. The risk was increased for MI too (OR = 2.7 (1.2–6.1)). Among women, the only significant association with IHD was for job dissatisfaction (OR = 3.0 (1.2–7.6)). **Conclusion:** In this population and in a period and society characterized by relative wealth and increasing employment rates, the Demand-control Model did not predict IHD. However, the feeling of job insecurity predicted both IHD and MI among men and job dissatisfaction predicted IHD among women.

Key words:

Ischemic heart disease, Myocardial infarction, Job strain, Psychological demands, Job satisfaction, Job insecurity

INTRODUCTION

During the last 30 years, epidemiological studies have provided growing evidence that work-related psychosocial factors play an important role in the development of ischemic heart disease (IHD) [1,2]. Most studies have been on the relationship between the Demand-Control

Model and IHD [3]. Several studies of cohorts in the 1970's and 80's found an increased risk for IHD among men who reported high psychological demands and low degree of control in their job [4–7]. In women similar findings have not been reported [8–10]. The early studies showed an association between control and IHD [11,12], while more

Address reprint request to B. Netterstrøm, Department of Occupational and Environmental Medicine, Bispebjerg University Hospital, Bispebjerg Bakke 23, 2400 Copenhagen NV (e-mail: bnet002@bbh.regionh.dk).

This Study complies with the Declaration of Helsinki. The research protocol of the Copenhagen City Heart Study has been approved by the locally appointed ethics committee.

recent studies found association between high demands and IHD [13,14]. Use of job title as an exposure assessment in US and Sweden has been conducted without significant results [12,15–17]. Recent studies found no association between either demands or control or the combination of these and IHD [2,18]. In a recent review we concluded that the found associations could fully be explained by the association between demands and disease risk. In addition we found insufficient evidence for a relationship between IHD and effort-reward imbalance, injustice, job insecurity or long working hours [2].

As the labour market demands have changed, there is a possibility that the perceived psychological strains which formerly were associated with IHD have changed too, which might result in that the Demand-Control Model is no longer a reliable tool to measure psychosocial risk factors for IHD [19]. Therefore, in this study we have chosen to deal also with other measures in analysing data from a cohort of employed urban citizens in relation to ischemic heart disease. Data from baseline has been reported earlier [20].

Participants

The study was carried out within the framework the Copenhagen City Heart Study [21]. The third round in the Copenhagen City Study was conducted from January 1, 1992 to June 13 1994. Participants were 63% of all invited persons living in an area of central Copenhagen. Those who participated in the period January 1993 to June 1994, aged 30 or more and at the same time were occupationally active were included in the study. In total, 595 women and 551 men were included. Their basic characteristics are shown in Table 1.

METHODS

All participants filled in a self-administered questionnaire regarding health and working conditions and went through a medical examination including measurements of height, weight, lung function, ECG and blood pressure. Fasting blood samples were tested for total serum chole-

sterol, high-density lipoprotein (HDL), plasma fibrinogen and plasma glucose.

All participants were traced in the Danish Hospitalization Registry and Death Registry from baseline till July 2007, which gives a follow up period between 13 and 14.5 years.

Independent variables

The two dimensions in the Demand-control Model were measured via a series of questions in the questionnaire. The demand dimension was measured with five questions, three of which came from the Whitehall II Study [22]. These three deal with speed, intensity, and time pressure at work (see Table 2). In addition, two questions were asked, one on normal speed of work and the other on alertness. The control dimension was measured with 13 questions, all from the Whitehall II Study, split into decision latitude and skill discretion.

Social relations were elucidated with questions about isolated work and about opportunity to speak to colleagues, while support was measured with four questions about support from colleagues and superiors. The questionnaire included questions on job satisfaction, responsibility and job insecurity as well (Table 2). Job insecurity was measured with a four-item scale.

Social status was defined as suggested by the National Institute of Social Research by using the coded answers in the questionnaire on job title and occupational sector:

Group 1: Self-employed with more than 20 employees, academics and white collar workers with more than 50 subordinates.

Group 2: Self-employed with 6–20 employees and white collar workers with 11–50 subordinates. Furthermore, self-employed and white collar workers with more than 2 years of formal education.

Group 3: self-employed with zero to 5 employees and white collar workers with 1 to 10 subordinates, Furthermore, self-employed and white collar workers with 1 to 2 years of formal education.

Group 4: White collar workers without subordinates and skilled workers.

Group 5: Unskilled workers.

Table 1. Basic characteristics for 595 women and 551 men in the Copenhagen City Heart Study who participated in the work-related data collection January 1993 to June 1994

Variable	Women (%)	Men (%)
Age groups (years)		
30–39	21.8	24.3
40–49	33.3	32.1
50–59	43.4	41.2
60–67	1.5	2.4
Civil status		
Married	64.7	75.9
Unmarried	13.8	15.1
Divorced	16.3	8.2
Widow	5.2	0.9
Social status		
Academics, Leader with > 50 subordinates, self-employed: > 20 subord.	15.6	21.1
White collar: 11–50 subord., self-employed 6–20 subord.	26.8	16.8
White collar 1–10 subord., self-employed 0–5 subord.	24.6	20.4
Skilled workers, white collar without subordinates	19.4	30.5
Unskilled workers	13.6	11.2
Weekly working hours		
< 29	14.7	4.4
30–37	54.1	41.7
38–49	29.4	28.9
> 50	5.4	18.7
Leader	25.2	33.4
Size of work place (employed, n)		
< 50	49.0	46.8
50–500	31.7	34.3
> 500	19.3	18.9
Family income (US\$)		
< 35.000	18.3	4.2
35.000–70.000	43.2	46.3
> 70.000	38.5	49.5
Physical activity in leisure time		
Passive	7.7	9.4
Light physical exercise 2–4 hours/week	55.8	45.6
Heavy physical exercise 2–4 hours/week or more	36.4	45.0
Smokers	46.7	47.7
Physical measures	Mean (SD)	Mean (SD)
BMI (kg/m ²)	24.4 (4.1)	26.0 (3.7)
Systolic Blood Pressure (mmHg)	126.5 (16.4)	135.2 (16.4)

SD — standard deviation.

Table 2. Odds ratio (OR) for first time hospitalisation or death due to IHD during 13 years follow up for work related exposures

Variable	Women OR (95% CI)	Men OR (95% CI)
Social status		
Group 2 / group 1 reference	0.9 (0.1–5.3)	2.2 (0.8–6.4)
Group 3 / group 1 reference	2.8 (0.9–8.9)	2.5 (1.2–5.2)
Group 4 / group 1 reference	4.0 (1.7–9.7)	1.2 (0.6–2.1)
Group 5 / group 1 reference	2.7 (1.1–6.5)	2.6 (1.3–5.1)
Family income (US\$)		
< 35.000 > 70.000	3.6 (0.4–29.9)	1.1 (0.4–2.9)
35.000–70.000 / > 70.000	4.6 (0.6–36.0)	1.3 (0.6–3.0)
Work organisation		
Leader	1.1 (0.5–2.4)	1.5 (0.9–2.5)
Shift work	0.4 (0.1–1.7)	0.8 (0.4–1.6)
Private/public sector	0.7 (0.3–1.4)	1.6 (0.9–2.7)
Weekly working hours		
30–37 / < 30	1.5 (0.5–4.6)	1.1 (0.3–3.8)
38–50 / < 30	1.0 (0.3–3.8)	1.0 (0.3–3.7)
> 50 / < 30	0.8 (0.1–7.8)	0.7 (0.2–2.7)
Psychosocial factors at work		
Demands		
How is your work pace? Too high/suitable, low	1.4 (0.4–5.3)	0.9 (0.4–2.1)
The work needs all attention and concentration: \geq 3/4 of the time/rarely	0.6 (0.3–1.3)	1.2 (0.6–2.2)
Time enough to do everything, Sometimes or rarely/often	1.0 (0.5–2.2)	1.2 (0.7–2.5)
You need to work fast; Often/sometimes or rarely	0.7 (0.3–1.5)	0.7 (0.4–1.3)
Necessary to work very concentrated; Often/sometimes or rarely	0.9 (0.2–3.2)	1.1 (0.5–2.3)
Decision authority		
Influence on what you do Often/sometimes or rarely	0.5 (0.2–1.1)	0.7 (0.5–1.3)
Is it necessary to do things over and over again: Sometimes or rarely/often	1.1 (0.3–3.9)	0.6 (0.3–1.2)
Can you choose between different ways to perform your work: Often/sometimes or rarely	1.2 (0.3–4.9)	0.6 (0.2–1.6)
I have something to say on decisions: Often/sometimes, rarely	0.9 (0.3–3.2)	0.8 (0.3–1.7)
Others makes decision about my work: Sometimes, rarely/often	1.5 (0.2–3.2)	0.5 (0.2–1.6)
Influence on work pace: Often/sometimes, rarely	0.8 (0.2–3.0)	0.7 (0.3–1.6)
Working time flexible: Often/sometimes, rarely	0.7 (0.1–3.4)	0.5 (0.2–1.3)
Influence on who I am working with: Often/sometimes, rarely	0.7 (0.3–5.1)	0.8 (0.5–2.0)
Influence on the work environment: Often/sometimes, rarely	1.6 (0.4–5.9)	0.7 (0.3–1.6)
I decide when to take a break: Often/sometimes, rarely	1.2 (0.7–2.5)	0.9 (0.5–1.6)
Skill discretion		
Possibilities to learn new things: Often/sometimes, rarely	1.5 (0.7–6.9)	0.6 (0.3–1.4)
Does your work need expertise: Often/sometimes, rarely	0.6 (0.3–1.3)	1.2 (0.7–2.0)
Do you have to take initiative: Often/sometimes, rarely	1.6 (0.7–3.1)	0.7 (0.4–1.2)
Social relations		
Work isolated: \geq 3/4 of the time / less	1.3 (0.6–2.8)	1.0 (0.6–1.7)
Have possibility to talk to colleagues during work: \geq 3/4 of the time / less	1.1 (0.5–2.2)	1.2 (0.7–1.9)

Table 2. Odds ratio (OR) for first time hospitalisation or death due to IHD during 13 years follow up for work related exposures — cont.

Variable	Women OR (95% CI)	Men OR (95% CI)
Social support		
How often do you get help and support from colleagues: Often/sometimes, rarely	0.4 (0.1–1.1)	1.0 (0.5–1.9)
How often are your colleagues willing to listen to your problems with work: Often/sometimes, rarely	0.5 (0.3–1.0)	0.8 (0.5–1.4)
How often do you get help and support from your nearest superior: Often/sometimes, rarely	0.7 (0.3–1.7)	0.8 (0.4–1.7)
How often are your nearest superior willing to listen to your problems with work: Often/sometimes, rarely	0.9 (0.4–1.8)	1.0 (0.6–1.7)
Job satisfaction		
Thought of seeking another job, Yes/no	0.8 (0.3–2.1)	0.6 (0.2–1.2)
Are you satisfied with your work: No, to a lesser degree / yes, very much, to some extent	3.8 (1.5–9.4)	0.7 (0.2–2.4)
Insecurity		
Are you worried for that		
You become unemployed: Yes/no	1.0 (0.5–2.7)	1.0 (0.6–1.7)
You are transferred to other job: Yes/no	2.0 (0.9–4.3)	0.7 (0.3–1.6)
You become superfluous due to new technology: Yes/no	1.4 (0.5–3.7)	1.8 (0.9–3.6)
You have difficulties to find a new job if unemployed with the qualifications you have: Yes/no	1.7 (0.8–3.4)	2.4 (1.4–4.1)
Responsibility		
Do you have sufficient responsibility in your work: Too much / yes, too little	0.9 (0.1–6.6)	1.5 (0.5–5.3)
Risk for health or costs for you or others if you make mistakes: 3/4 of the time / less	0.9 (0.4–1.9)	0.6 (0.4–1.1)

Statistical analyses

Of the 1,146 participants, 70 men and 34 women contracted IHD during the follow up period (ICD 8 410–414, ICD 10 I20–I25). This outcome was used in the analyses, and no distinction was made between death and hospitalisation due to IHD. Furthermore, the 38 cases of MI (ICD 8 410, ICD 10 I20) among men and 11 among women were analysed in relation to the significant independent variables found in the former analyses. All analyses were carried out for each gender separately.

In the statistical analyses, we used the Chi²-test for discrete variables and Student's t-test for continuous variables in bivariate analyses. Adjustment for confounders was carried out using logistic regression analyses, where the adjustment factors were forced into the model.

RESULTS

Table 2 gives an overview of the statistical associations between work-related factors and the incidence of IHD during follow up. Social status had a significant association to

outcome as expected in both genders. In women, significantly increased risk was found among workers compared to group 1. Among men, an excess risk was found among unskilled workers and group 3 consisting of small enterprise employers and white collar workers with few subordinates or low degree of education. Family income showed no significant association with IHD, nor did weekly working hours, shift work or employment in the private sector. None of the items on psychological demands showed any significant association with IHD nor did any of the items concerning decision authority or skill discretion. In men, all associations with the items on decision authority were in the expected direction, but still insignificant. In women, the pattern was mixed. Even so, no significant associations between IHD and decision latitude were found for men (OR = 0.7 (0.4–1.2)) even after adjustment for age. However, among men more than 50 years of age, high demands predicted IHD significantly (OR = 2.3 (1.1–4.8)) which led to the significant associations between high strain or belonging to the active group and IHD. Social relations and social support were not associated significantly to IHD.

However, in women, social support from colleagues was borderline associated.

Job dissatisfaction was significantly associated with IHD in women, but not in men. Even though job dissatisfaction was linked to “thoughts of seeking another job”, this last item did not show any association to IHD.

In men, concern about low chances to find a new job if unemployed, given one’s current qualifications, was significantly associated with IHD, while the concern about unemployment was not. The associations between IHD and the psychosocial factors at work are shown in Table 3. After adjustment for social status and coronary risk factors, no significant associations were found except for insecurity among men and low job satisfaction among women. The positive association between IHD and insecurity among men held for MI even after adjustment for all confounders (OR = 3.2 (1.4–7.1)). Due to lack of MI cases among women with low job satisfaction, no relevant statistical tests could be made concerning MI. Among men above 50, high demands were positively associated with IHD, which resulted in significant associations

for IHD for both the active and strained group in the Demand-Control Model.

As social status was the most prominent confounder, the data were split into three strata of social status in order to get a more differentiated picture of the results.

The tendency to increased risk for Job strain was in men almost U-shaped, as there was no excess risk in group 3. It were, however, demands which contributed to the high risk estimate for strain in group 1 and 2 and control in group 4 and 5. Among women, no particular patterns showed. Significantly elevated risk was only found for workers (group 4 and 5), in women for low job satisfaction (OR = 3.2 (1.1–10.0)) and among men for insecurity (OR = 2.7 (1.2–6.3)).

Discussion

This cohort study of employed urban citizens showed no support for the opinion that the Demand Control Model is valuable as a measure of psychosocial risk factors in the work environment for the development of IHD. This is in line with another Danish study conducted 8 years earlier

Table 3. Odds ratio OR for first time hospitalisation or death due to IHD during follow up for exposures in the psychosocial work environment

Psychosocial factor	Adjustment for age OR (95% CI)		Adjustment for age, social status and coronary risk factors OR (95% CI)					
	women	men	< 50 years		> 50 years		All	
			women	men	women	men	women	men
Demands, high/low	0.9 (0.4–1.8)	1.1 (0.6–1.8)	1.1 (0.3–4.2)	0.7 (0.3–1.7)	0.9 (0.3–2.4)	2.3 (1.1–4.8)	1.1 (0.5–1.6)	1.5 (0.8–2.6)
Control, high/low	0.6 (0.3–1.4)	0.7 (0.4–1.1)	1.9 (0.4–9.9)	1.3 (0.5–3.4)	0.9 (0.3–2.7)	0.7 (0.3–1.6)	1.2 (0.5–3.1)	0.9 (0.5–1.3)
Job strain								
Active	1.3 (0.7–2.2)	1.2 (0.8–1.9)	0.8 (0.1–6.2)	0.6 (0.2–2.0)	(0.2–6.1)	3.2 (1.1–9.0)	1.1 (0.2–4.7)	1.5 (0.5–3.5)
Passive	0.8 (0.4–1.6)	0.9 (0.5–1.4)	0.2 (0.01–1.8)	0.7 (0.2–2.3)	1.3 (0.2–7.7)	1.8 (0.6–5.6)	0.8 (0.2–3.6)	1.1 (0.5–2.7)
Strain	1.2 (0.6–2.3)	1.3 (0.8–2.1)	0.6 (0.1–3.4)	0.5 (0.1–1.0)	1.1 (0.2–6.0)	3.5 (1.1–10.5)	1.1 (0.3–4.2)	1.6 (0.7–3.7)
Social support, high/low	0.6 (0.3–1.2)	0.8 (0.5–1.4)	1.0 (0.3–3.4)	1.0 (0.4–2.3)	0.7 (0.3–1.8)	0.7 (0.4–1.5)	0.9 (0.4–1.4)	0.8 (0.5–1.4)
Social relations, many/few	1.2 (0.6–3.3)	1.6 (0.9–2.5)	1.7 (0.5–5.9)	1.6 (0.7–3.7)	0.8 (0.3–2.2)	1.5 (0.7–2.9)	1.1 (0.5–2.0)	1.6 (0.9–2.7)
Conflicts	0.5 (0.3–1.1)	1.2 (0.7–2.0)	0.8 (0.2–2.7)	1.0 (0.4–2.6)	0.6 (0.2–1.7)	1.2 (0.6–2.4)	0.7 (0.2–3.2)	1.1 (0.6–3.1)
Insecurity, high/low	1.6 (0.7–3.4)	2.1 (1.2–3.6)	1.1 (0.3–4.3)	1.3 (0.5–3.3)	1.6 (0.6–4.3)	2.7 (1.2–5.7)	1.4 (0.5–3.2)	2.5 (1.1–5.6)
Job satisfaction, low/high	3.7 (1.5–9.1)	0.9 (0.7–3.2)	0.8 (0.1–5.4)	1.8 (0.4–6.9)	4.0 (1.2–13.1)	+	3.0 (1.2–7.6)	0.9 (0.3–3.3)

+ No cases in the low job satisfaction group.

in the suburb of Copenhagen, where we found a clear association between self-reported high psychological demands at work and IHD independently of other coronary risk factors, but no association between the combination of demands and control and IHD [14]. Other recent studies have also failed to support the Demand Control Model hypothesis regarding IHD [18,24], while older studies have shown good predicting of IHD using the model [1,2]. The difference in those findings may be due to the changes in exposures to psychosocial risk factors during the time elapsed between the early and recent studies. In Denmark, the changes in the labour market during the last decade of the last century have resulted in longer working hours, more computerised and highly skilled work and a decrease in numbers of workers with clerical, cleaning and industrial jobs [19]. The study has some intrinsic methodological limitations, as do other similar studies, and this might have affected the results. Firstly, exposure was only measured at baseline, and the relatively long follow-up time means that it must be assumed that a certain number of participants have stopped working and consequently have not been exposed for some time. Further, the exposure intensity might have changed during the follow-up time, and these factors add to an underestimation of the observed associations.

The fact that only 63% of the entire study population participated in the study constitutes a selection bias. We do not know anything about the non-participants' health or working conditions, we only know that their age and gender distribution compared to that of the participants. It is difficult to imagine that information bias could influence the results in any systematic way. The Danish hospitalisation register has a very high degree of validity, and the procedures used in Denmark with regard to diagnosing IHD are considered to be of a high standard [23]. This means that it is unlikely that there are cases of misclassification when it comes to the endpoints.

The confounder control could lead to overcontrol, but the risk estimates were mostly affected by control for social status and not for conventional risk factors.

Consequently, this cohort study partly supports the findings from other studies indicating that job insecurity could

be a major risk factor for IHD [25–27]. This is of importance in the light of the financial crisis which did not affect the results of this study. Further studies are needed in order to determine whether these are isolated findings or perhaps indications of a new trend due to changing working conditions at the modern labour market.

The results also indicate that risk for IHD associated with psychosocial factors at work is unevenly distributed in the labour force.

REFERENCES

1. Kivimäki M, Virtanen M, Elovainio M, et al. *Work stress in the etiology of coronary heart disease — A meta-analysis*. Scand J Work Environ Health 2006;32:431–42.
2. Eller NH, Netterstrøm B, Gyntelberg F, Kristensen TS, Nielsen F, Steptoe A, Theorell T. *Work-related psychosocial factors and the development of ischemic heart disease. A systematic review*. Cardiol Rev 2009;17:83–97.
3. Belkic KL, Landsbergis PA, Schnall PL, et al. *Is job strain a major source of cardiovascular disease risk?* Scand J Work Environ Health 2004;30:85–128.
4. Haan M. *Job strain and cardiovascular disease: A ten years prospective study*. Am J Epidemiol 1985;122:532–4.
5. Netterstrom B, Nielsen FE, Kristensen TS, Bach E, Moller L. *Relation between job strain and myocardial infarction: A case-control study*. Occup Environ Med 1999;56:339–42.
6. Kivimäki M, Leino-Arjas P, Luukkonen R, Riihimäki H, Vahtera J, Kirjonen J. *Work stress and risk of cardiovascular mortality: Prospective cohort study of industrial employees*. BMJ 2002;325:857–60.
7. Kivimäki M, Ferrie JE, Brunner E, Head J, Shipley MJ, Vahtera J, et al. *Justice at work and reduced risk of coronary heart disease among employees: The Whitehall II Study*. Arch Intern Med 2005;165:2245–51.
8. Kuper H, Marmot M. *Job strain, job demands, decision latitude, and risk of coronary heart disease within the Whitehall II study*. J Epidemiol Community Health 2003;57:147–53.
9. Andre-Petersson L, Engstrom G, Hedblad B, Janzon L, Rosvall M. *Social support at work and the risk of myocardial infarction and stroke in women and men*. Soc Sci Med 2007;64: 830–41.

10. Eaker ED, Sullivan LM, Kelly-Hayes M, D'Agostino RB Sr., Benjamin EJ. *Does job strain increase the risk for coronary heart disease or death in men and women? The Framingham Offspring Study.* Am J Epidemiol 2004;159:950–8.
11. Alfredsson L, Karasek R, Theorell T. *Myocardial infarction risk and the psychosocial work environment: an analysis of the male Swedish working force.* Soc Sci Med 1982;16:463–7.
12. Steenland K, Johnson J, Nowlin S. *A follow-up study of job strain and heart disease among males in the NHANES I population.* Am J Ind Med 1997;31:256–60.
13. Kornitzer M, De Smet P, Sans S, Dramaix M, Boulenguez C, DeBacker G, et al. *Job stress and major coronary events: results from the Job Stress, Absenteeism and Coronary Heart Disease in Europe study.* Eur J Cardiovasc Prev Rehabil 2006;13:695–704.
14. Netterstrom B, Kristensen TS, Sjol A. *Psychological job demands increase the risk of ischaemic heart disease: a 14-year cohort study of employed Danish men.* Eur J Cardiovasc Prev Rehabil 2006;13:414–20.
15. Hammar N, Alfredsson L, Johnson JV. *Job strain, social support at work, and incidence of myocardial infarction.* Occup Environ Med 1998;55:548–53.
16. Alterman T, Shekelle RB, Vernon SW, Bureau KD. *Decision latitude, psychological demand, job strain and coronary heart disease in the Western Electric Study.* Am J Epidemiol 1994;139:620–7.
17. Reed DM, LaCroix AZ, Karasek RA, Miller D, MacLean CA. *Occupational strain and the incidence of coronary heart disease.* Am J Epidemiol 1989;129:495–502.
18. De Bacquer D, Pelfrene E, Clays E, Mak R, Moreau M, de Smet P. *Perceived job stress and incidence of coronary events: 3-year follow-up of the Belgian Job Stress Project cohort.* Am J Epidemiol 2005;161:434–41.
19. Burr H, Bjørner J, Kristensen TS, Tüchsen F, Bach E. *Trends in the Danish work environment 1999–2000 and their relation to labor force changes.* Scand J Work Environ Health 2003;29:270–9.
20. Netterstrøm B, Kristensen TS, Møller L, Jensen G, Schnohr P. *Angina pectoris, job strain and social status.* Int J Behav Med 1998;5:312–22.
21. Appleyard M, Hansen A, Schnohr P, Jensen G, Nyboe J. *The Copenhagen City Heart Study.* Scand J Soc Med 1987; Suppl 41:1–160.
22. North FM, Syme SL, Feeney A, Shipley M, Marmot M. *Psychosocial work environment and sickness absence among British civil servants: The Whitehall II study.* Am J Public Health 1996;86:332–40.
23. Tüchsen F, Andersen O, Olsen J. *Referral bias among health workers in studies using hospitalization as a proxy of the underlying incidence rate.* J Clin Epidemiol 1996;7:791–4.
24. Bonde JP, Munch-Hansen T, Agerbo E, Suadicani P, Wiclaw J, Westergaard-Nielsen N. *Job strain and ischemic heart disease: A prospective study using a new approach for exposure assessment.* JOM 2009;51:732–8.
25. Iversen L, Sabroe S, Damsgaard MT. *Hospital admissions before and after shipyard closure.* BMJ 1989;299:1073–6.
26. Ferrie JE, Shipley MJ, Marmot MG, Stansfeld S, Davey Smith G. *The health effects of major organisational change and job insecurity.* Soc Sci Med 1998;46:243–54.
27. Vahtera J, Kivimaki M, Pentti J, Linna A, Virtanen M, Virtanen P, et al. *Organisational downsizing, sickness absence, and mortality: 10-town prospective cohort study.* BMJ 2004;328:555. DOI 10.1136/bmj.37972.496262.0D.